

Fuzzy Set - Basics

Prof. Dr. Rudolf Kruse

Motivation

Every day humans use imprecise linguistic terms
e.g. big, fast, about 12 o'clock, old, etc.

All complex human actions are decisions based on such concepts:

- driving and parking a car,
- financial/business decisions,
- law and justice,
-

So, these terms and the way they are processed play a crucial role for the development of intelligent systems .

Computers need a mathematical model to express and process such complex semantics.

Concepts of classical mathematics are often inadequate for such models.

Example – The Sorites Paradox

If a pile of sand is **small**, adding one grain of sand to it leaves it **small**.

A pile of sand with a single grain is **small**.

Hence all sand dunes are **small**.

Paradox comes from an all-or-nothing treatment of *small*.

It is better to say that the degree of truth of the statement “pile of sand is small” decreases by adding one grain after another.

Example - Solution of the Sorites Paradox

Statement $A(n)$: “ n grains of sand form a sand pile.”

Let $d_n = T(A(n))$ denote “degree of truth” for $A(n)$. Then

$$0 = d_0 \leq d_1 \leq \dots \leq d_n \leq \dots \leq 1$$

can be seen as truth values of a **many valued logic**

with 0 (false) and 1 (true).

The concept of a Fuzzy Logic is useful in such cases

(not everything is either black or white, often there are degrees or grey-levels)

Example - Imprecise Linguistic Expressions

Consider the notion *bald*:

A person without hair on his head is bald,
a very hairy person is definitely not bald.


Usually, *bald* is only partly applicable to a person.

A *baldness/non baldness* threshold is counter-intuitive?

Solution: Use „generalized sets“ with membership degrees.

bald?

truth degree



The concept of a Fuzzy set is useful in such cases.

Lotfi Asker Zadeh

Classes of objects in the real world do not have precisely defined criteria of membership.

Such imprecisely defined “classes” play an important role in human thinking and the development of intelligent systems.

Particularly in domains of pattern recognition, communication of information, and abstraction.



Lotfi A. Zadeh's Principle of Incompatibility

“Stated informally, the essence of this principle is that as the complexity of a system increases, our ability to make precise and yet significant statements about its behavior diminishes until a threshold is reached beyond which precision and significance (or relevance) become almost mutually exclusive characteristics.”

Fuzzy sets/fuzzy logic are used as mechanism for abstraction of unnecessary or too complex details.

Applications using Fuzzy Sets

Control Engineering

Approximate Reasoning



Data Sciences

Artificial Intelligence



Rudolf Kruse received IEEE Fuzzy Pioneer Award for „Learning Methods for Fuzzy Systems“ in 2018

Fuzzy sets as generalizations of classical sets

ling. description		model
all numbers smaller than 10	\longrightarrow <i>objective</i>	 characteristic function of a set
all numbers <u>almost</u> equal to 10	\longrightarrow <i>subjective</i>	 membership function of a "fuzzy set"

Definition

A fuzzy set μ of X is a function from the reference set X to the unit interval, *i.e.* $\mu : X \rightarrow [0, 1]$. $F(X)$ represents the set of all fuzzy sets of X , *i.e.* $F(X) := \{\mu \mid \mu : X \rightarrow [0, 1]\}$.

Membership Functions


$\mu_M(u) = 1$ reflects full membership in M .

$\mu_M(u) = 0$ expresses absolute non-membership in M .

Sets can be viewed as special case of fuzzy sets where only full membership and absolute non-membership are allowed.

Such sets are called *crisp sets* or Boolean sets.

Membership degrees $0 < \mu_M < 1$ represent *partial membership*.



Representing *young* in "a young person"

Membership Function


A Membership function attached to a given linguistic description (such as *young*) depends on the context – it is subjective.

A **young** retired person is certainly older than a **young** student.


Membership degrees are fixed only *by convention*:

Unit interval as range of membership grades is arbitrary but easy to use.

Examples: Characteristic Function of a Set




exactly two




between 1 and 3

Examples: Fuzzy Sets are described by Functions



Approximately 2




Approximately 2

Exact numerical value has membership degree of 1.

Left: monotonically increasing, right: monotonically decreasing,
i.e. unimodal function.

Example – Velocity of Rotating Hard Disk



Fuzzy set μ characterizing the **normal** velocity of rotating hard disk.

Let v be the velocity of rotating hard disk in revolutions per minute.

Modelling of expert's knowledge:

“It's *impossible* that v drops under a or exceeds d .


It's highly certain that any value between $[b, c]$ can occur.”

Otherwise I defined my subjective point of view , I also used data“


Vertical Representation

So far, fuzzy sets were described by their characteristic/membership function and assigning degree of membership $\mu(x)$ to each element $x \in X$.

That is the **vertical representation** of the corresponding fuzzy set, e.g. linguistic expression like “about m ”



or “approximately between b and c ”




Cuts of Fuzzy sets


Let $\mu \in \mathcal{F}(X)$ and $\alpha \in [0, 1]$. Then the sets

$$[\mu]_{\alpha} = \{x \in X \mid \mu(x) \geq \alpha\}, \quad [\mu]_{\underline{\alpha}} = \{x \in X \mid \mu(x) > \alpha\}$$

are called the α -cut and *strict* α -cut of μ .



An Example




Let μ be triangular function on \mathbb{R} as shown above.

α -cut of μ can be constructed by

1. drawing horizontal line parallel to x-axis through point $(0, \alpha)$,
2. projecting this section onto x-axis.

$$[\mu]_\alpha = \begin{cases} [a + \alpha(m - a), b - \alpha(b - m)], & \text{if } 0 < \alpha \leq 1, \\ \mathbb{R}, & \text{if } \alpha = 0. \end{cases}$$

Properties of α -cuts I



Theorem


Let $\mu \in \mathcal{F}(X)$, $\alpha \in [0, 1]$ and $\beta \in [0, 1]$.

(a) $[\mu]_0 = X$,

(b) $\alpha < \beta \implies [\mu]_\alpha \supseteq [\mu]_\beta$,

(c) $\bigcap_{\alpha: \alpha < \beta} [\mu]_\alpha = [\mu]_\beta$.

Characteristic function of a set



Let $A \subseteq X$, $\chi_A : X \rightarrow [0, 1]$

$$\chi_A(x) = \begin{cases} 1 & \text{if } x \in A, \\ 0 & \text{otherwise} \end{cases}$$

Then $[\chi_A]_\alpha = A$ for $0 < \alpha \leq 1$.

χ_A is called indicator function or characteristic function of A .

Theorem (Representation Theorem)


Let $\mu \in \mathcal{F}(X)$. Then

$$\mu(x) = \sup_{\alpha \in [0,1]} \left\{ \min(\alpha, \chi_{[\mu]_\alpha}(x)) \right\}$$


where $\chi_{[\mu]_\alpha}(x) = \begin{cases} 1, & \text{if } x \in [\mu]_\alpha \\ 0, & \text{otherwise.} \end{cases}$

So, fuzzy set can be obtained as upper envelope of its α -cuts.


Simply draw α -cuts parallel to horizontal axis in height of α .



Vertical View



Horizontal View




Example: Modelling (Approximately 5) or (Greater than or Equal to 7)

Suppose that $X = [0, 15]$.

An expert chooses $L = \{0, 0.25, 0.5, 0.75, 1\}$ and α -cuts:

- $A_0 = [0, 15]$,
- $A_{0.25} = [3, 15]$,
- $A_{0.5} = [4, 6] \cup [7, 15]$,
- $A_{0.75} = [4.5, 5.5] \cup [7, 15]$,
- $A_1 = \{5\} \cup [7, 15]$.




The family $(A_\alpha)_{\alpha \in L}$ of sets induces upper shown fuzzy set.

“Approximately 5 or greater than or equal to 7”

μ_A is obtained as upper envelope of the family A of sets.


The difference between horizontal and vertical view is obvious:



The horizontal representation is easier to process in computers.

Also, restricting the domain of x-axis to a discrete set is usually done.

Horizontal Representation in the Computer



Fuzzy sets are usually stored as chain of linear lists.

A finite union of closed intervals is stored by their bounds.

This data structure is appropriate for arithmetic operators.

Fuzzy Set Representation as Systems of Cuts

In this manner we obtain **system of sets**

$$\mathcal{A} = (A_\alpha)_{\alpha \in L}, \quad L \subseteq [0, 1], \quad \text{card}(L) \in \mathbb{N}.$$

\mathcal{A} must satisfy consistency conditions for $\alpha, \beta \in L$:

- (a) $0 \in L \implies A_0 = X$, (fixing of reference set)
- (b) $\alpha < \beta \implies A_\alpha \supseteq A_\beta$. (monotonicity)



This induces fuzzy set

$$\begin{aligned} \mu_{\mathcal{A}} : X &\rightarrow [0, 1], \\ \mu_{\mathcal{A}}(x) &= \sup_{\alpha \in L} \{ \min(\alpha, \chi_{A_\alpha}(x)) \}. \end{aligned}$$

If L is not finite but comprises all values $[0, 1]$, then μ must satisfy

- (c) $\bigcap_{\alpha: \alpha < \beta} A_\alpha = A_\beta$. (condition for continuity)

Convex Fuzzy Sets are easy to handle



A fuzzy set $\mu \in F(\mathbb{R})$ is convex if and only if

$$\mu(\lambda x_1 + (1 - \lambda)x_2) \geq \min\{\mu(x_1), \mu(x_2)\}$$

for all $x_1, x_2 \in \mathbb{R}$ and all $\lambda \in [0, 1]$.